

LINEAR ALGEBRA

and Learning

from Data

First Edition

MANUAL FOR INSTRUCTORS

Gilbert Strang

Massachusetts Institute of Technology

math.mit.edu/weborder.php (orders)

math.mit.edu/~gs

www.wellesleycambridge.com

email: linearalgebrabook@gmail.com

Wellesley - Cambridge Press

Box 812060

Wellesley, Massachusetts 02482

Problem Set I.1, page 6

- 1 A combination of \mathbf{u} , \mathbf{v} , and $\mathbf{u} + \mathbf{v}$ (vectors in \mathbf{R}^4) produces

$$\mathbf{u} + \mathbf{v} - (\mathbf{u} + \mathbf{v}) = \mathbf{0} \quad \left[\begin{array}{ccc} \mathbf{u} & \mathbf{v} & \mathbf{u} + \mathbf{v} \end{array} \right] \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \mathbf{0} \quad A\mathbf{x} = \mathbf{0}$$

A is 4 by 3, \mathbf{x} is 3 by 1, $\mathbf{0}$ is 4 by 1. Your example could use numerical vectors.

- 2 Suppose $A\mathbf{x} = A\mathbf{y}$. Then if $\mathbf{z} = c(\mathbf{x} - \mathbf{y})$ for any number c , we have $A\mathbf{z} = \mathbf{0}$. One candidate is always the zero vector $\mathbf{z} = \mathbf{0}$ (from the choice $c = 0$).
- 3 We are given vectors \mathbf{a}_1 to \mathbf{a}_n in \mathbf{R}^m with $c_1\mathbf{a}_1 + \cdots + c_n\mathbf{a}_n = \mathbf{0}$.

(1) At the matrix level $A\mathbf{c} = \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \mathbf{c} = \mathbf{0}$, with the \mathbf{a} 's in the columns of A , and c 's in the vector \mathbf{c} .

(2) At the scalar level this is $\sum_{j=1}^n a_{ij}c_j = 0$ for each row $i = 1, 2, \dots, m$ of A .

- 4 Two vectors \mathbf{x} and \mathbf{y} out of many solutions to $A\mathbf{x} = \mathbf{0}$ for $A = \mathbf{ones}(3, 3)$ are

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

These vectors $\mathbf{x} = (1, 1, -2)$ and $\mathbf{y} = (3, -3, 0)$ are independent. But there is no 3rd vector \mathbf{z} with $A\mathbf{z} = \mathbf{0}$ and independent $\mathbf{x}, \mathbf{y}, \mathbf{z}$. (If there were, then combinations of $\mathbf{x}, \mathbf{y}, \mathbf{z}$ would say that every vector \mathbf{w} solves $A\mathbf{w} = \mathbf{0}$, which is not true.)

- 5 (a) The vector $\mathbf{z} = (1, -1, 1)$ is perpendicular to $\mathbf{v} = (1, 1, 0)$ and $\mathbf{w} = (0, 1, 1)$. Then \mathbf{z} is perpendicular to all combinations of \mathbf{v} and \mathbf{w} —a whole plane in \mathbf{R}^3 .
- (b) $\mathbf{u} = (1, 1, 1)$ is NOT a combination of \mathbf{v} and \mathbf{w} . And \mathbf{u} is NOT perpendicular to $\mathbf{z} = (1, -1, 1)$: Their dot product is $\mathbf{u}^T \mathbf{z} = 1$.

6 If u, v, w are corners of a parallelogram, then $z = \text{corner 4}$ can be $u + v - w$ or $u - v + w$ or $-u + v + w$. Here those 4th corners are $z = (4, 0)$ or $z = (-2, 2)$ or $z = (4, 4)$.

Reasoning: The corners A, B, C, D around a parallelogram have $A + C = B + D$.

7 The column space of $A = \begin{bmatrix} v & w & v + 2w \end{bmatrix}$ consists of all combinations of v and w .

Case 1 v and w are independent. Then $\mathbf{C}(A)$ has dimension 2 (a *plane*). A has rank 2 and its nullspace is a line (dimension 1) in \mathbf{R}^3 : Then $2 + 1 = 3$.

Case 2 w is a multiple cv (not both zero). Then $\mathbf{C}(A)$ is a line and the nullspace is a plane: $1 + 2 = 3$.

Case 3 $v = w = 0$ and the nullspace of A (= zero matrix) is all of \mathbf{R}^3 : $0 + 3 = 3$.

$$\mathbf{8} \quad A = \begin{bmatrix} 1 & 4 & 9 \\ 1 & 4 & 9 \\ 1 & 4 & 9 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 9 \end{bmatrix} = \text{rank-1 matrix.}$$

9 If $\mathbf{C}(A) = \mathbf{R}^3$ then $m = 3$ and $n \geq 3$ and $r = 3$.

$$\mathbf{10} \quad A_1 = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 9 & -6 \\ 2 & 6 & -4 \end{bmatrix} \text{ has } C_1 = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} \quad A_2 = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \text{ has } C_2 = \begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 7 & 8 \end{bmatrix}$$

$$\mathbf{11} \quad A_1 = C_1 R_1 = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 3 & -2 \end{bmatrix} \quad A_2 = C_2 R_2 = \begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

12 The vector $(1, 3, 2)$ is a basis for $\mathbf{C}(A_1)$. The vectors $(1, 4, 7)$ and $(2, 5, 8)$ are a basis for $\mathbf{C}(A_2)$. The dimensions are 1 and 2, so the ranks of the matrices are 1 and 2. Then A_1 and A_2 must have 1 and 2 independent rows.

$$\mathbf{13} \quad \text{An example is } A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix}. \text{ Then } C \text{ and } R \text{ are } 4 \text{ by } 2 \text{ and } 2 \text{ by } 4.$$

14 $A = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 2 \\ 0 & 2 \end{bmatrix}$ or $B = \begin{bmatrix} 0 & 2 & 5 \\ 0 & 2 & 5 \end{bmatrix}$ have the same column

spaces but different row spaces. The basic columns chosen directly from A and B are $(1, 1)$ and $(2, 2)$. The rank = number of vectors in the column basis must be the same (1).

15 If $A = CR$, then the numbers in row 1 of C multiply the rows of R to produce row 1 of A .

16 “The rows of R are a basis for the row space of A ” means: R has independent rows, and every row of A is a combination of the rows of R .

17 $A_1 = \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} = C_1 R_1$ $A_2 = \begin{bmatrix} C_1 \\ C_1 \end{bmatrix} \begin{bmatrix} R_1 \end{bmatrix}$ $A_3 = \begin{bmatrix} C_1 \\ C_1 \end{bmatrix} \begin{bmatrix} R_1 & R_1 \end{bmatrix}$

18 If $A = CR$ then $\begin{bmatrix} 0 & A \\ 0 & A \end{bmatrix} = \begin{bmatrix} C \\ C \end{bmatrix} \begin{bmatrix} 0 & R \end{bmatrix}$

19 $A = \begin{bmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 3 & 8 \\ 0 & -1 & -2 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 \\ 0 & -1 & -2 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 \\ 0 & -1 & -2 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow$
 $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} = \text{same row space as } A.$ Remove the zero row to see R in $A = CR$.

20 $C = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ gives $C^T C = \begin{bmatrix} 13 \end{bmatrix}$. $R = \begin{bmatrix} 2 & 4 \end{bmatrix}$ produces $RR^T = \begin{bmatrix} 20 \end{bmatrix}$.

$A = \begin{bmatrix} 2 & 4 \\ 3 & 6 \end{bmatrix}$ produces $C^T A R^T = \begin{bmatrix} 130 \end{bmatrix}$. Then $M = \frac{1}{13} \begin{bmatrix} 130 \end{bmatrix} \frac{1}{20} = \begin{bmatrix} \frac{1}{2} \end{bmatrix}$.

$$\mathbf{21} \quad \mathbf{C}^T \mathbf{C} = \begin{bmatrix} 1 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 5 \\ 5 & 14 \end{bmatrix} \text{ has } (\mathbf{C}^T \mathbf{C})^{-1} = \frac{1}{3} \begin{bmatrix} 14 & -5 \\ -5 & 2 \end{bmatrix}$$

$$\mathbf{R} \mathbf{R}^T = \begin{bmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 3 & 2 \\ 8 & 6 \end{bmatrix} = \begin{bmatrix} 74 & 55 \\ 55 & 41 \end{bmatrix} \text{ has } (\mathbf{R} \mathbf{R}^T)^{-1} = \frac{1}{9} \begin{bmatrix} 41 & -55 \\ -55 & 74 \end{bmatrix}$$

$$\mathbf{C}^T \mathbf{A} \mathbf{R}^T = \begin{bmatrix} 1 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 3 & 2 \\ 8 & 6 \end{bmatrix} = \begin{bmatrix} 2 & 5 & 14 \\ 5 & 14 & 38 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 3 & 2 \\ 8 & 6 \end{bmatrix} =$$

$$\begin{bmatrix} 129 & 96 \\ 351 & 261 \end{bmatrix}$$

$$\mathbf{M} = \frac{1}{3} \begin{bmatrix} 14 & -5 \\ -5 & 2 \end{bmatrix} \begin{bmatrix} 129 & 96 \\ 351 & 261 \end{bmatrix} \frac{1}{9} \begin{bmatrix} 41 & -55 \\ -55 & 74 \end{bmatrix} = ?$$

$$\mathbf{22} \quad \text{If } \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & ma \\ c & mc \end{bmatrix} \text{ then } ad - bc = mac - mac = 0: \text{ dependent columns!}$$

$$\mathbf{23} \quad \begin{bmatrix} 2 & 4 \\ 2 & 4 \\ 2 & 4 \end{bmatrix} = \mathbf{C} \mathbf{R} = \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} \begin{bmatrix} \frac{1}{2} \end{bmatrix} \begin{bmatrix} 2 & 4 \end{bmatrix} = \mathbf{C} \mathbf{M} \mathbf{R}$$

(row of \mathbf{R} from \mathbf{A})

$$\mathbf{24} \quad \begin{bmatrix} 2 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix} = \mathbf{C} \mathbf{R} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \mathbf{C} \mathbf{M} \mathbf{R}$$

Problem Set I.2, page 13

- 1 $A \begin{bmatrix} \mathbf{x} & \mathbf{y} \end{bmatrix} = \begin{bmatrix} \mathbf{0} & \mathbf{0} \end{bmatrix}$ where $B = \begin{bmatrix} \mathbf{x} & \mathbf{y} \end{bmatrix}$ is n by 2 and $C = \begin{bmatrix} \mathbf{0} & \mathbf{0} \end{bmatrix}$ is m by 2.
- 2 Yes, $\mathbf{a}\mathbf{b}^T$ is an m by n matrix. The number $a_i b_j$ is in row i , column j of $\mathbf{a}\mathbf{b}^T$. If $\mathbf{b} = \mathbf{a}$ then $\mathbf{a}\mathbf{a}^T$ is a *symmetric* matrix.
- 3 (a) $AB = \mathbf{a}_1 \mathbf{b}_1^T + \cdots + \mathbf{a}_n \mathbf{b}_n^T$
 (b) The i, j entry of AB is $\sum_{k=1}^n a_{ik} b_{kj}$.
- 4 If B has one column $\begin{bmatrix} b_1 & \cdots & b_n \end{bmatrix}^T$ then $AB = \mathbf{a}_1 b_1 + \cdots + \mathbf{a}_n b_n =$ combination of the columns of A (as expected). Each row of B is one number b_k .
- 5 Verify $(AB)C = A(BC)$ for $AB = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} = \begin{bmatrix} b_1 + ab_3 & b_2 + ab_4 \\ b_3 & b_4 \end{bmatrix}$
 and $BC = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ c & 1 \end{bmatrix} = \begin{bmatrix} b_1 + cb_2 & b_2 \\ b_3 + cb_4 & b_4 \end{bmatrix}$ AB was row ops
 BC was col ops
 Row ops then col ops $\begin{bmatrix} b_1 + ab_3 & b_2 + ab_4 \\ b_3 & b_4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ c & 1 \end{bmatrix} = \begin{bmatrix} b_1 + ab_3 + cb_2 + acb_4 & b_2 + ab_4 \\ b_3 + cb_4 & b_4 \end{bmatrix}$
 Col ops then row ops $\begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} b_1 + cb_2 & b_2 \\ b_3 + cb_4 & b_4 \end{bmatrix} = \begin{bmatrix} b_1 + ab_3 + cb_2 + acb_4 & b_2 + ab_4 \\ b_3 + cb_4 & b_4 \end{bmatrix}$ **SAME**
 If A, C were both row operations, $(AC)B = (CA)B$ would usually be false.
- 6 $B = I$ has rows $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3 = (1, 0, 0), (0, 1, 0), (0, 0, 1)$. The rank-1 matrices are
 $\mathbf{a}_1 \mathbf{b}_1 = \begin{bmatrix} \mathbf{a}_1 & \mathbf{0} & \mathbf{0} \end{bmatrix}$ $\mathbf{a}_2 \mathbf{b}_2 = \begin{bmatrix} \mathbf{0} & \mathbf{a}_2 & \mathbf{0} \end{bmatrix}$ $\mathbf{a}_3 \mathbf{b}_3 = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{a}_3 \end{bmatrix}$.
 The sum of those rank-1 matrices is $AI = A$.
- 7 If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ then $AB = \begin{bmatrix} 2 & 0 \\ 4 & 0 \end{bmatrix}$ has a smaller column space than A . Note that $(\text{row space of } AB) \leq (\text{row space of } B)$.
- 8 For $k = 1$ to n
 For $i = 1$ to m
 For $j = 1$ to p

Problem Set I.3, page 20

1 If $B\mathbf{x} = \mathbf{0}$ then $AB\mathbf{x} = \mathbf{0}$. So every \mathbf{x} in the nullspace of B is also in the nullspace of AB .

2 $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ has $A^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and the rank has dropped.

But $A^T A = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ has the same nullspace and rank as A .

3 If $C = \begin{bmatrix} A \\ B \end{bmatrix}$ then $C\mathbf{x} = \mathbf{0}$ requires both $A\mathbf{x} = \mathbf{0}$ and $B\mathbf{x} = \mathbf{0}$. So the nullspace of C is the **intersection** $\mathbf{N}(A) \cap \mathbf{N}(B)$.

4 Actually row space = column space requires nullspace of $A =$ nullspace of A^T . But it *does not* require symmetry. Choose any invertible matrix like $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$.

5 $r = m = n$ A_1 is any invertible square matrix

$r = m < n$ A_2 has extra columns like $A_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$

$r = n < m$ A_3 has extra rows like A_2^T


$r < m, r < n$ $A_4 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ or $\begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}$ or $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}$.

6 First, if $A\mathbf{x} = \mathbf{0}$ then $A^T A\mathbf{x} = \mathbf{0}$. So $\mathbf{N}(A^T A)$ contains (or equals) $\mathbf{N}(A)$.

Second, if $A^T A\mathbf{x} = \mathbf{0}$ then $\mathbf{x}^T A^T A\mathbf{x} = 0$ and $\|A\mathbf{x}\|^2 = 0$. Then $A\mathbf{x} = \mathbf{0}$ and $\mathbf{N}(A)$ contains (or equals) $\mathbf{N}(A^T A)$. Altogether $\mathbf{N}(A^T A)$ **equals** $\mathbf{N}(A)$.

7 $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $A^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ have *different* nullspaces.

8 $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ has $\mathbf{C}(A) = \mathbf{N}(A) =$ all vectors $\begin{bmatrix} x_1 \\ 0 \end{bmatrix}$. But $\mathbf{C}(A) = \mathbf{N}(A^T)$ is impossible.

9  **8 edges**
5 nodes

$Ax = 0$ for $x = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$

Columns 1 to 5 are *dependent*
Columns 1 to 4 are *independent*

Incidence matrix $A = \begin{bmatrix} -1 & 1 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix}$

has rank 4
 $\mathbf{N}(A)$ has dimension 1
 $\mathbf{N}(A^T)$ has dimension 4
 $8 - 4 = 4$ small loops

10 If $\mathbf{N}(A) = \{\mathbf{0}\}$, the nullspace of $B = \begin{bmatrix} A & A & A \end{bmatrix}$ contains all vectors $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ with $x + y + z = \mathbf{0}$.

- 11 (i) $S \cap T$ has dimension 0, 1, or 2
(ii) $S + T$ has dimension 7, 8, or 9
(iii) $S^\perp =$ (vectors perpendicular to S) has dimension $10 - 2 = 8$.

Problem Set I.4, page 27

$$\begin{aligned}
 \mathbf{1} \quad & \begin{bmatrix} 2 & 1 \\ 6 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 0 & 4 \end{bmatrix} \\
 & \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\
 & \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1/2 & 1 & 0 \\ 0 & -2/3 & 1 \end{bmatrix} \begin{bmatrix} 2 & -1 & 0 \\ 0 & 3/2 & -1 \\ 0 & 0 & 4/3 \end{bmatrix}
 \end{aligned}$$

$$\mathbf{2} \quad a_{ij} = a_{i1}a_{1j}/a_{11} \quad \text{Check } A = \begin{bmatrix} 2 & 3 \\ 4 & 6 \end{bmatrix} \text{ and } a_{22} = (4)(3)/(2).$$

If $a_{11} = 0$ then the formula breaks down. We could still have rank 1.

$$\mathbf{3} \quad EA = U \text{ is } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 4 & 2 \\ 6 & 3 & 5 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 4 & 2 \\ 0 & 0 & 5 \end{bmatrix}$$

$$A = LU \text{ is } \begin{bmatrix} 2 & 1 & 0 \\ 0 & 4 & 2 \\ 6 & 3 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 4 & 2 \\ 0 & 0 & 5 \end{bmatrix} \quad \text{Note } L = E^{-1}$$

$$\mathbf{4} \quad E_2E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -c & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ -b & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ac-b & -c & 1 \end{bmatrix} \quad \begin{array}{l} ac - b \text{ mixes} \\ \text{the multipliers} \end{array}$$

$$E_1^{-1}E_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & c & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & c & 1 \end{bmatrix} \quad \begin{array}{l} \text{In this order the} \\ \text{multipliers fall} \\ \text{into place in } L \end{array}$$

- 5** If zero appears in a pivot position then $A = LU$ is **not possible**. We need a *permutation* P to exchange rows and lead to nonzero pivots.

$$\begin{bmatrix} 0 & 1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \ell & 1 \end{bmatrix} \begin{bmatrix} d & e \\ 0 & f \end{bmatrix} \text{ leads to } \begin{array}{l} 0 = d \quad (1, 1 \text{ entry}) \\ 2 = \ell d \quad (\text{impossible if } d = 0) \end{array}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \ell & 1 & 0 \\ m & n & 1 \end{bmatrix} \begin{bmatrix} d & e & g \\ 0 & f & h \\ 0 & 0 & i \end{bmatrix} \text{ leads to } \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \ell & 1 \end{bmatrix} \begin{bmatrix} d & e \\ 0 & f \end{bmatrix}$$

second pivot is zero

$$\text{Then } \begin{bmatrix} d & e \\ 0 & f \end{bmatrix} \text{ must be singular and } \begin{bmatrix} d & e & g \\ 0 & f & h \\ 0 & 0 & i \end{bmatrix} \text{ is singular. BUT } \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix}$$

is invertible! So $A = LU$ is again impossible.

- 6** $c = 2$ makes the second pivot zero. But A is still invertible.

$c = 1$ makes the third pivot zero. Then A is singular.

$$\mathbf{7} \quad A = LU \text{ is } \begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & 0 & d-c \end{bmatrix}$$

For nonzero pivots in U , we need $a \neq 0, b \neq a, c \neq b, d \neq c$.

- 8** If A is tridiagonal and $A = LU$ (no row exchanges in elimination) then L and U have *two diagonals*. The only elimination steps subtract a pivot row from the row directly beneath it.

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & & \\ 1 & 1 & \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ & 1 & 1 \\ & & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} a & a & 0 \\ a & a+b & b \\ 0 & b & b+c \end{bmatrix} = \begin{bmatrix} 1 & & \\ 1 & 1 & \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & 0 \\ & b & b \\ & & c \end{bmatrix}$$

- 9** The second pivot in elimination depends only on the upper left 2 by 2 submatrix A_2 of A . The third pivot depends only on the upper left 3 by 3 submatrix (and so on). So if the pivots (diagonal entries in U) are 5, 9, 3, then the pivots for A_2 are 5, 9.

- 10** Continuing Problem 9, the upper left parts of L and U **come from the upper left part of A** . Then $L_k U_k$ is the factorization of A_k .

$$A = \begin{bmatrix} L_k & 0 \\ * & * \end{bmatrix} \begin{bmatrix} U_k & * \\ 0 & * \end{bmatrix} \text{ so } A_k = L_k U_k$$

- 11** The example could exchange rows of A to put the larger number 3 into the (1,1) position where it would become the first pivot. That would be the usual permutation in MATLAB and other systems.

This problem also exchanges columns to put the even larger number 4 into the (1,1) position. A column exchange comes from a permutation multiplying on the *right side* of A . So this problem works on both sides :

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \text{ leads to } P_1 A P_2 = \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \text{ leads to } P_2 A P_1 = \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix}$$

- 12** With m rows and n columns and $m < n$, elimination normally leads from A to

$$U = \begin{bmatrix} U_1 & U_2 \\ m \times n & m \times (n-m) \end{bmatrix} \quad \text{Example: } U = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 9 \end{bmatrix}$$

There must be nonzero solutions to $U\mathbf{x} = \mathbf{0}$. To see this, set $x_3 = 1$ and solve

$$\begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = - \begin{bmatrix} 4 \\ 9 \end{bmatrix} \text{ to find } \begin{matrix} x_1 = 2 \\ x_2 = -3 \end{matrix}. \text{ So } \mathbf{x} = \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} \text{ solves } A\mathbf{x} = \mathbf{0}.$$

Problem Set I.5, page 35

1 If $\mathbf{u}^T \mathbf{v} = 0$ and $\mathbf{u}^T \mathbf{u} = 1$ and $\mathbf{v}^T \mathbf{v} = 1$, then $(\mathbf{u} + \mathbf{v})^T (\mathbf{u} - \mathbf{v}) = 1 + 0 - 0 - 1 = \mathbf{0}$

Also $\|\mathbf{u} + \mathbf{v}\|^2 = \mathbf{u}^T \mathbf{u} + \mathbf{v}^T \mathbf{u} + \mathbf{u}^T \mathbf{v} + \mathbf{v}^T \mathbf{v} = 1 + 0 + 0 + 1 = 2$ and $\|\mathbf{u} - \mathbf{v}\|^2 = 2$

2 \mathbf{v} is separated into a piece $\mathbf{u}(\mathbf{u}^T \mathbf{v})$ in the direction of \mathbf{u} and the remaining piece $\mathbf{w} = \mathbf{v} - \mathbf{u}(\mathbf{u}^T \mathbf{v})$ perpendicular to \mathbf{u} . Check $\mathbf{u}^T \mathbf{w} = \mathbf{u}^T \mathbf{v} - (\mathbf{u}^T \mathbf{u})(\mathbf{u}^T \mathbf{v}) = 0$.

3 $\mathbf{w}^T \mathbf{w} + \mathbf{z}^T \mathbf{z} = (\mathbf{u} + \mathbf{v})^T (\mathbf{u} + \mathbf{v}) + (\mathbf{u} - \mathbf{v})^T (\mathbf{u} - \mathbf{v}) = (\mathbf{u}^T \mathbf{u} + \mathbf{v}^T \mathbf{v} + \mathbf{u}^T \mathbf{v} + \mathbf{v}^T \mathbf{u}) + (\mathbf{u}^T \mathbf{u} + \mathbf{v}^T \mathbf{v} - \mathbf{u}^T \mathbf{v} - \mathbf{v}^T \mathbf{u}) = 2(\mathbf{u}^T \mathbf{u} + \mathbf{v}^T \mathbf{v})$.

Sum of squares of 2 diagonals = Sum of squares of 4 sides.

4 Check $(Q\mathbf{x})^T (Q\mathbf{y}) = \mathbf{x}^T Q^T Q \mathbf{y} = \mathbf{x}^T \mathbf{y}$: Angles are preserved when all vectors are multiplied by Q . Remember $\mathbf{x}^T \mathbf{y} = \|\mathbf{x}\| \|\mathbf{y}\| \cos \theta = (Q\mathbf{x})^T (Q\mathbf{y})$: same θ !

5 If Q is orthogonal (this word assumes a square matrix) then $Q^T Q = I$ and Q^T is Q^{-1} . Check $(Q_1 Q_2)^T = Q_2^T Q_1^T = Q_2^{-1} Q_1^{-1}$ which is $(Q_1 Q_2)^{-1}$.

6 Every permutation matrix has unit vectors in its columns (single 1 and $n - 1$ zeros). Those columns are orthogonal because their 1's are in different positions.

$$7 \quad PF = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & i^2 & i^3 \\ 1 & i^2 & i^4 & i^6 \\ 1 & i^3 & i^6 & i^9 \end{bmatrix} = \begin{bmatrix} 1 & i & i^2 & i^3 \\ 1 & i^2 & i^4 & i^6 \\ 1 & i^3 & i^6 & i^9 \\ 1 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & i^2 & i^3 \\ 1 & i^2 & i^4 & i^6 \\ 1 & i^3 & i^6 & i^9 \end{bmatrix} \begin{bmatrix} 1 \\ i \\ i^2 \\ i^3 \end{bmatrix}$$

This says that P times the 4 columns of F gives those same 4 columns times $1, i, i^2, i^3 = \lambda_1, \lambda_2, \lambda_3, \lambda_4 =$ the 4 eigenvalues of P .

The columns of $F/2$ are orthonormal! To check, remember that for the dot product of two *complex vectors*, we take complex conjugates of the first vector: *change i to $-i$* .

8 $W^T W = \begin{bmatrix} 4 & & & \\ & 4 & & \\ & & 2 & \\ & & & 2 \end{bmatrix}$ so that the columns of W are orthogonal but not orthonormal.

Then $W^{-1} = (W^T W)^{-1} W^T = \begin{bmatrix} 1/4 & & & \\ & 1/4 & & \\ & & 1/2 & \\ & & & 1/2 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$.

Problem Set I.6, page 41

- 1** To check: $\lambda_1 + \lambda_2 = \text{trace} = \cos \theta + \cos \theta$ and $\lambda_1 \lambda_2 = \text{determinant} = \cos^2 \theta + \sin^2 \theta = 1$ and $\bar{x}_1^T x_2 = 0$ (orthogonal matrices have complex orthogonal eigenvectors).

$$Q^{-1} = Q^T \text{ has eigenvalues } \frac{1}{e^{i\theta}} = e^{-i\theta} \text{ and } \frac{1}{e^{-i\theta}} = e^{i\theta}$$

- 2** $\det \begin{bmatrix} -\lambda & 2 \\ 1 & 1 - \lambda \end{bmatrix} = \lambda^2 - \lambda - 2 = (\lambda - 2)(\lambda + 1) = 0$ gives $\lambda_1 = 2$ and $\lambda_2 = -1$.

The sum $2 - 1$ agrees with the trace $0 + 1$. A^{-1} has the same eigenvectors as A , with eigenvalues $\lambda_1^{-1} = \frac{1}{2}$ and $\lambda_2^{-1} = -1$.

- 3** A has $\lambda = 3$ and 1 , B has $\lambda = 1$ and 3 , $A + B$ has $\lambda = 5$ and 3 . Eigenvalues of $A + B$ are generally **not equal** to $\lambda(A) + \lambda(B)$. Now A and B have $\lambda = 1$ (*repeated*). AB and BA both have $\lambda^2 - 4\lambda + 1 = 0$ (leading to $\lambda = 2 \pm \sqrt{3}$ by the quadratic formula). The eigenvalues of AB and BA are the same—but not equal to $\lambda(A)$ times $\lambda(B)$.

- 4** A and B have $\lambda_1 = 1$ and $\lambda_2 = 1$. AB and BA have $\lambda^2 - 4\lambda + 1$ and the quadratic formula gives $\lambda = 2 \pm \sqrt{3}$. Eigenvalues of AB are *not equal* to eigenvalues of A times eigenvalues of B . Eigenvalues of AB and BA **are equal** (this is proved at the end of Section 6.2).

- 5** (a) Multiply Ax to see λx which reveals λ (b) Solve $(A - \lambda I)x = 0$ to find x .

- 6** $\det(A - \lambda I) = \lambda^2 - 1.4\lambda + 0.4$ so A has $\lambda_1 = 1$ and $\lambda_2 = 0.4$ with $x_1 = (1, 2)$ and $x_2 = (1, -1)$. A^∞ has $\lambda_1 = 1$ and $\lambda_2 = 0$ (same eigenvectors). A^{100} has $\lambda_1 = 1$ and $\lambda_2 = (0.4)^{100}$ which is near zero. So A^{100} is very near A^∞ : same eigenvectors and close eigenvalues.

- 7** Set $\lambda = 0$ in $\det(A - \lambda I) = (\lambda_1 - \lambda) \dots (\lambda_n - \lambda)$ to find $\det A = (\lambda_1)(\lambda_2) \dots (\lambda_n)$.

- 8** $\lambda_1 = \frac{1}{2}(a + d + \sqrt{(a - d)^2 - 4bc})$ and $\lambda_2 = \frac{1}{2}(a + d - \sqrt{(a - d)^2 - 4bc})$ add to $a + d$. If A has $\lambda_1 = 3$ and $\lambda_2 = 4$ then $\det(A - \lambda I) = (\lambda - 3)(\lambda - 4) = \lambda^2 - 7\lambda + 12$.

- 9** These 3 matrices have $\lambda = 4$ and 5 , trace 9 , det 20 : $\begin{bmatrix} 4 & 0 \\ 0 & 5 \end{bmatrix}$, $\begin{bmatrix} 3 & 2 \\ -1 & 6 \end{bmatrix}$, $\begin{bmatrix} 2 & 2 \\ -3 & 7 \end{bmatrix}$.

- 10** $A = \begin{bmatrix} 0 & 1 \\ -28 & 11 \end{bmatrix}$ has trace 11 and determinant 28, so $\lambda = 4$ and 7. Moving to a 3 by 3 companion matrix, for eigenvalues 1, 2, 3 we want $\det(C - \lambda I) = (1 - \lambda)(2 - \lambda)(3 - \lambda)$. Multiply out to get $-\lambda^3 + 6\lambda^2 - 11\lambda + 6$. To get those numbers 6, -11, 6 from a companion matrix you just put them into the last row:

$$C = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix} \text{ Notice the trace } 6 = 1 + 2 + 3 \text{ and determinant } 6 = (1)(2)(3).$$

- 11** $(A - \lambda I)$ has the same determinant as $(A - \lambda I)^T$ because every square matrix has $\det M = \det M^T$. Pick $M = A - \lambda I$.

$$\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \text{ and } \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \text{ have different eigenvectors.}$$

- 12** $\lambda = 0, 0, 6$ (notice rank 1 and trace 6). Two eigenvectors of uv^T are perpendicular to v and the third eigenvector is u : $x_1 = (0, -2, 1)$, $x_2 = (1, -2, 0)$, $x_3 = (1, 2, 1)$.

- 13** When A and B have the same n λ 's and x 's, look at any combination $v = c_1x_1 + \dots + c_nx_n$. Multiply by A and B : $Av = c_1\lambda_1x_1 + \dots + c_n\lambda_nx_n$ **equals** $Bv = c_1\lambda_1x_1 + \dots + c_n\lambda_nx_n$ **for all vectors** v . So $A = B$.

- 14** (a) u is a basis for the nullspace (we know $Au = 0u$); v and w give a basis for the column space (we know Av and Aw are in the column space).

(b) $A(v/3 + w/5) = 3v/3 + 5w/5 = v + w$. So $x = v/3 + w/5$ is a particular solution to $Ax = v + w$. Add any cu from the nullspace

(c) If $Ax = u$ had a solution, u would be in the column space: wrong dimension 3.

- 15** Eigenvectors in X and eigenvalues in Λ . Then $A = X\Lambda X^{-1}$ is given below.

The second matrix has $\lambda = 0$ (rank 1) and $\lambda = 4$ (trace = 4). A new $A = X\Lambda X^{-1}$:

$$\begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 3 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{bmatrix}.$$

16 If $A = X\Lambda X^{-1}$ then the eigenvalue matrix for $A + 2I$ is $\Lambda + 2I$ and the eigenvector matrix is still X . So $A + 2I = X(\Lambda + 2I)X^{-1} = X\Lambda X^{-1} + X(2I)X^{-1} = A + 2I$.

17 (a) False: We are not given the λ 's (b) True (c) True (d) False: For this we would need the eigenvectors of X .

$$\mathbf{18} \quad A = X\Lambda X^{-1} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \lambda_1 & \\ & \lambda_2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} / 2 = \begin{bmatrix} \lambda_1 + \lambda_2 & \lambda_1 - \lambda_2 \\ \lambda_1 - \lambda_2 & \lambda_1 + \lambda_2 \end{bmatrix} / 2.$$

These are the matrices $\begin{bmatrix} a & b \\ b & a \end{bmatrix}$, their eigenvectors are $(1, 1)$ and $(1, -1)$.

19 (a) *True* (no zero eigenvalues) (b) *False* (repeated $\lambda = 2$ may have only one line of eigenvectors) (c) *False* (repeated λ may have a full set of eigenvectors)

20 (a) False: don't know if $\lambda = 0$ or not.

(b) True: an eigenvector is missing, which can only happen for a repeated eigenvalue.

(c) True: We know there is only one line of eigenvectors.

21 $A^k = X\Lambda^k X^{-1}$ approaches zero **if and only if every** $|\lambda| < 1$; A_1 is a Markov matrix so $\lambda_{\max} = 1$ and $A_1^k \rightarrow A_1^\infty$, A_2 has $\lambda = .6 \pm .3$ so $A_2^k \rightarrow 0$.

$$\mathbf{22} \quad \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} = X\Lambda X^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \text{ and}$$

$$A^k = X\Lambda^k X^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3^k \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}.$$

$$\text{Multiply those last three matrices to get } A^k = \frac{1}{2} \begin{bmatrix} 1 + 3^k & 1 - 3^k \\ 1 - 3^k & 1 + 3^k \end{bmatrix}.$$

$$\mathbf{23} \quad R = X\sqrt{\Lambda}X^{-1} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 3 & \\ & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} / 2 = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \text{ has } R^2 = A.$$

\sqrt{B} needs $\lambda = \sqrt{9}$ and $\sqrt{-1}$, trace (their sum) is not real so \sqrt{B} cannot be real. Note that $\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$ has two imaginary eigenvalues $\sqrt{-1} = i$ and $-i$, real trace 0, real square root $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

24 $A = X\Lambda_1X^{-1}$ and $B = X\Lambda_2X^{-1}$. Diagonal matrices always give $\Lambda_1\Lambda_2 = \Lambda_2\Lambda_1$.

Then $AB = BA$ from

$$X\Lambda_1X^{-1}X\Lambda_2X^{-1} = X\Lambda_1\Lambda_2X^{-1} = X\Lambda_2\Lambda_1X^{-1} = X\Lambda_2X^{-1}X\Lambda_1X^{-1} = BA.$$

25 Multiply columns of X times rows of ΛX^{-1} .

26 To have $A = B\Lambda B^{-1}$ requires A to have a full set of n independent eigenvectors. Then B is the *eigenvector matrix* and it is invertible.

Problem Set I.7, page 52

- 1 The key is to form $\mathbf{y}^T \mathbf{S} \mathbf{x}$ in two ways, using $S^T = S$ to make them agree. One way starts with $S\mathbf{x} = \lambda\mathbf{x}$: multiply by \mathbf{y}^T . The other way starts with $S\mathbf{y} = \alpha\mathbf{y}$ and then $\mathbf{y}^T S^T = \alpha\mathbf{y}^T$.

The final step finds $0 = (\lambda - \alpha) \mathbf{y}^T \mathbf{x}$ which forces $\mathbf{y}^T \mathbf{x} = 0$.

- 2 Only $S_4 = \begin{bmatrix} 1 & 10 \\ 10 & 101 \end{bmatrix}$ has two positive eigenvalues since $101 > 10^2$.

$\mathbf{x}^T S_1 \mathbf{x} = 5x_1^2 + 12x_1x_2 + 7x_2^2$ is negative for example when $x_1 = 4$ and $x_2 = -3$:

A_1 is not positive definite as its determinant confirms; S_2 has trace c_0 ; S_3 has $\det = 0$.

- 3 Positive definite $\begin{bmatrix} 1 & 0 \\ b & 1 \end{bmatrix} \begin{bmatrix} 1 & b \\ 0 & 9 - b^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ b & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 9 - b^2 \end{bmatrix} \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix} = LDL^T$
for $-3 < b < 3$

Positive definite $\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 \\ 0 & c - 8 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & c - 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = LDL^T$
for $c > 8$

Positive definite $L = \begin{bmatrix} 1 & 0 \\ -b/c & 1 \end{bmatrix}$ $D = \begin{bmatrix} c & 0 \\ 0 & c - b/c \end{bmatrix}$ $S = LDL^T$
for $c^2 > b$

- 4 If \mathbf{x} is not real then $\lambda = \mathbf{x}^T A \mathbf{x} / \mathbf{x}^T \mathbf{x}$ is *not* always real. Can't assume real eigenvectors!

- 5 $\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} = 2 \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{bmatrix} + 4 \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}$; $\begin{bmatrix} 9 & 12 \\ 12 & 16 \end{bmatrix} = 0 \begin{bmatrix} .64 & -.48 \\ -.48 & .36 \end{bmatrix} + 25 \begin{bmatrix} .36 & .48 \\ .48 & .64 \end{bmatrix}$

- 6 M is skew-symmetric and **orthogonal**; λ 's must be $i, i, -i, -i$ to have trace 0, $|\lambda| = 1$.

- 7 $A = \begin{bmatrix} i & 1 \\ 1 & -i \end{bmatrix}$ has $\lambda = 0, 0$ and only one independent eigenvector $\mathbf{x} = (i, 1)$. The good property for complex matrices is not $A^T = A$ (symmetric) but $\overline{A}^T = A$ (Hermitian with real eigenvalues and orthogonal eigenvectors: see Problem 22 and Section 9.2).

- 8 Eigenvectors $(1, 0)$ and $(1, 1)$ give a 45° angle even with A^T very close to A .

9 (a) $S^T = S$ and $S^T S = I$ lead to $S^2 = I$.

(b) The only possible eigenvalues of S are 1 and -1 .

(c) $\Lambda = \begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix}$ so $S = \begin{bmatrix} Q_1 & Q_2 \end{bmatrix} \Lambda \begin{bmatrix} Q_1^T \\ Q_2^T \end{bmatrix} = Q_1 Q_1^T - Q_2 Q_2^T$ with $Q_1^T Q_2 = 0$.

10 Eigenvalues of $A^T S A$ are different from eigenvalues of S but the signs are the same: the Law of Inertia gives the same number of plus-minus-zero eigenvalues.

11 $\det(S - aI) = \begin{vmatrix} 0 & b \\ b & c - a \end{vmatrix} = -b^2$ is negative. So the point $x = a$ is between the two eigenvalues where $\det(S - \lambda_1 I) = 0$ and $\det(S - \lambda_2 I) = 0$. This $\lambda_1 \leq a \leq \lambda_2$ is a general rule for larger matrices too (Section II.2): Eigenvalues of the submatrix of size $n - 1$ **interlace** eigenvalues of the n by n symmetric matrix.

12 $\mathbf{x}^T S \mathbf{x} = 2x_1 x_2$ comes from $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$. That matrix has eigenvalues 1 and -1 . Conclusion: Saddle points are associated with eigenvalues of *both signs*.

13 $A^T A = \begin{bmatrix} 1 & 2 \\ 2 & 13 \end{bmatrix}$ and $A^T A = \begin{bmatrix} 6 & 5 \\ 5 & 6 \end{bmatrix}$ are positive definite; $A^T A = \begin{bmatrix} 2 & 3 & 3 \\ 3 & 5 & 4 \\ 3 & 4 & 5 \end{bmatrix}$ is singular (and positive semidefinite). The first two A 's have independent columns. The 2 by 3 A cannot have full column rank 3, with only 2 rows; $A^T A$ is singular.

14 $S = \begin{bmatrix} 4 & -4 & 8 \\ -4 & 4 & -8 \\ 8 & -8 & 16 \end{bmatrix}$ has only one pivot = 4, rank $S = 1$, eigenvalues are 24, 0, 0, $\det S = 0$.

15 Corner determinants $|S_1| = 2$, $|S_2| = 6$, $|S_3| = 30$. The pivots are $2/1$, $6/2$, $30/6$.

16 S is positive definite for $c > 1$; determinants c , $c^2 - 1$, and $(c - 1)^2(c + 2) > 0$.

T is *never* positive definite (determinants $d - 4$ and $-4d + 12$ are never both positive).

17 $S = \begin{bmatrix} 1 & 5 \\ 5 & 10 \end{bmatrix}$ is an example with $a + c > 2b$ but $ac < b^2$, so not positive definite.

18 $\mathbf{x}^T S \mathbf{x}$ is zero when $(x_1, x_2, x_3) = (0, 1, 0)$ because of the zero on the diagonal. Actually $\mathbf{x}^T S \mathbf{x}$ goes *negative* for $\mathbf{x} = (1, -10, 0)$ because the second pivot is *negative*.

- 19** If a_{jj} were smaller than all λ 's, $S - a_{jj}I$ would have all eigenvalues > 0 (positive definite). But $S - a_{jj}I$ has a zero in the (j, j) position; impossible by Problem 18.

$$\mathbf{20} \quad A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \sqrt{9} & \\ & \sqrt{1} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}; \quad A = Q \begin{bmatrix} 4 & 0 \\ 0 & 2 \end{bmatrix} Q^T = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}.$$

- 21** The ellipse $x^2 + xy + y^2 = 1$ has axes with half-lengths $1/\sqrt{\lambda} = \sqrt{2}$ and $\sqrt{2/3}$.

$$\mathbf{22} \quad \text{The Cholesky factors } A = (L\sqrt{D})^T = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \text{ and } A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & \sqrt{5} \end{bmatrix} \text{ have}$$

square roots of the pivots from D . Note again $A^T A = LDL^T = S$.

- 23** The energy test gives $\mathbf{x}^T(A^T C A) \mathbf{x} = (A\mathbf{x})^T C (A\mathbf{x}) = \mathbf{y}^T C \mathbf{y} > 0$ since C is positive definite and $\mathbf{y} = A\mathbf{x}$ is only zero if \mathbf{x} is zero. (A was assumed to have independent columns.)

This is just like the $A^T A$ discussion, but now with a positive definite C in $A^T C A$.

$$\mathbf{24} \quad S_1 = \begin{bmatrix} 6x^2 & 2x \\ 2x & 2 \end{bmatrix} \text{ is semidefinite; } f_1 = (\frac{1}{2}x^2 + y)^2 = 0 \text{ on the curve } \frac{1}{2}x^2 + y = 0;$$

$$S_2 = \begin{bmatrix} 6x & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \text{ is indefinite at } (0, 1) \text{ where first derivatives} = 0. \text{ Then}$$

$x = 0, y = 1$ is a saddle point of the function $f_2(x, y)$.

- 25** If $c > 9$ the graph of z is a bowl, if $c < 9$ the graph has a saddle point. When $c = 9$ the graph of $z = (2x + 3y)^2$ is a "trough" staying at zero along the line $2x + 3y = 0$.

- 26** $\det S = (1)(10)(1) = 10$; $\lambda = 2$ and 5 ; $\mathbf{x}_1 = (\cos \theta, \sin \theta)$, $\mathbf{x}_2 = (-\sin \theta, \cos \theta)$; the λ 's are positive. So S is positive definite.

- 27** Energy $\mathbf{x}^T S \mathbf{x} = a(x_1 + x_2 + x_3)^2 + c(x_2 - x_3)^2 \geq 0$ if $a \geq 0$ and $c \geq 0$: semidefinite.

S has rank ≤ 2 and determinant = 0; cannot be positive definite for any a and c .

- 28** (a) The eigenvalues of $\lambda_1 I - S$ are $\lambda_1 - \lambda_1, \lambda_1 - \lambda_2, \dots, \lambda_1 - \lambda_n$. Those are ≥ 0 ; $\lambda_1 I - S$ is semidefinite.
- (b) Semidefinite matrices have energy $\mathbf{x}^T (\lambda_1 I - S) \mathbf{x} \geq 0$. Then $\lambda_1 \mathbf{x}^T \mathbf{x} \geq \mathbf{x}^T S \mathbf{x}$.
- (c) Part (b) says $\mathbf{x}^T S \mathbf{x} / \mathbf{x}^T \mathbf{x} \leq \lambda_1$ for all \mathbf{x} . Equality holds at the leading eigenvector with $S \mathbf{x} = \lambda_1 \mathbf{x}$.

(Note that the maximum is λ_1 —the first printing missed the subscript “one”).

Problem Set I.8, page 68

1 $(c_1 \mathbf{v}_1^T + \cdots + c_n \mathbf{v}_n^T)(c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n) = c_1^2 + \cdots + c_n^2$ because the \mathbf{v} 's are orthonormal.

$$\begin{aligned} (c_1 \mathbf{v}_1^T + \cdots + c_n \mathbf{v}_n^T) S (c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n) &= (\quad) (c_1 \lambda_1 \mathbf{v}_1 + \cdots + c_n \lambda_n \mathbf{v}_n) \\ &= c_1^2 \lambda_1 + \cdots + c_n^2 \lambda_n. \end{aligned}$$

2 Remember that $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$. Then $\lambda_1 c_1^2 + \cdots + \lambda_n c_n^2 \leq (c_1^2 + \cdots + c_n^2)$.

Therefore the ratio $R(\mathbf{x})$ is $\leq \lambda_1$. It equals λ_1 when $\mathbf{x} = \mathbf{v}_1$.

3 Notice that $\mathbf{x}^T \mathbf{v}_1 = (c_1 \mathbf{v}_1^T + \cdots + c_n \mathbf{v}_n^T) \mathbf{v}_1 = c_1$. Then $\mathbf{x}^T \mathbf{v}_1 = 0$ means $c_1 = 0$.

Now $R(\mathbf{x}) = \frac{\lambda_2 c_2^2 + \cdots + \lambda_n c_n^2}{c_2^2 + \cdots + c_n^2}$ is a maximum when $\mathbf{x} = \mathbf{v}_2$ and $c_2 = 1$ and other c 's = 0.

4 The maximum of $R(\mathbf{x}) = \mathbf{x}^T S \mathbf{x} / \mathbf{x}^T \mathbf{x}$ is λ_3 when \mathbf{x} is restricted by $\mathbf{x}^T \mathbf{v}_1 = \mathbf{x}^T \mathbf{v}_2 = 0$.

5 If $A = U \Sigma V^T$ then $A^T = V \Sigma^T U^T$ (singular vectors \mathbf{u} and \mathbf{v} are reversed but the numbers $\sigma_1, \dots, \sigma_r$ do not change. Then $A \mathbf{v} = \sigma \mathbf{u}$ and $A^T \mathbf{u} = \sigma \mathbf{v}$ for each pair of singular vectors.

For example $A = \begin{bmatrix} 0 & 5 \\ 0 & 0 \end{bmatrix}$ has $\sigma_1 = 5$ and so does $A^T = \begin{bmatrix} 0 & 0 \\ 5 & 0 \end{bmatrix}$. But $\|A \mathbf{x}\| \neq \|A^T \mathbf{x}\|$ for most \mathbf{x} .

6 Exchange \mathbf{u} 's and \mathbf{v} 's (and keep $\sigma = \sqrt{45}$ and $\sigma = \sqrt{5}$) in equation (12) = the SVD of $\begin{bmatrix} 3 & 0 \\ 4 & 5 \end{bmatrix}$.

7 This question should have told us which matrix norm to use! If we use $\|A\| = \sigma_1$ then removing $\sigma_1 \mathbf{u}_1 \mathbf{v}_1^T$ will leave the norm as σ_2 . If we use the Frobenius norm $(\sigma_1^2 + \cdots + \sigma_r^2)^{1/2}$, then removing $\sigma_1 \mathbf{u}_1 \mathbf{v}_1^T$ will leave $(\sigma_2^2 + \cdots + \sigma_r^2)^{1/2}$.

$$\mathbf{8} \quad \begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = U \Sigma V^T$$

- 9** (*Correction to first printing*) Remove both factors $\frac{1}{2}$ that multiply $\mathbf{x}^T S \mathbf{x}$. Then maximizing $\mathbf{x}^T S \mathbf{x}$ with $\mathbf{x}^T \mathbf{x} = 1$ is the same as maximizing their ratio $R(\mathbf{x})$.

Now the gradient of $L = \mathbf{x}^T S \mathbf{x} + \lambda(\mathbf{x}^T \mathbf{x} - 1)$ is $2S\mathbf{x} - 2\lambda\mathbf{x}$. This gives *gradient* = 0 at all eigenvectors \mathbf{v}_1 to \mathbf{v}_n . Testing $R(\mathbf{x})$ at each eigenvector gives $R(\mathbf{v}_k) = \lambda_k$ so $\mathbf{x} = \mathbf{v}_1$ maximizes $R(\mathbf{x})$.

- 10** If you remove columns of a matrix, this cannot increase the norm. Reason: We still have $\text{norm} = \max \|A\mathbf{v}\|/\|\mathbf{v}\|$ but we are only keeping the \mathbf{v} 's with zeros in the positions corresponding to removed columns. So the maximum can only move down and never up.

Then removing columns of the transpose (rows of the original matrix) can only reduce the norm further. So a submatrix of A cannot have larger norm than A .

- 11** The trace of $S = \begin{bmatrix} 0 & A & \\ & & A^T & 0 \end{bmatrix}$ is zero. The eigenvalues of S come in plus-minus pairs so they add to zero. If $A = \text{diag}(1, 2, \dots, n)$ is diagonal, these $2n$ eigenvalues of S are 1 to n and -1 to $-n$. The $2n$ eigenvectors of S have 1 in positions 1 to n with all $+1$ or all -1 in positions $n+1$ to $2n$.

12 $A = \begin{bmatrix} 2 & 4 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix}$ means that $A = \frac{\begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}}{\sqrt{5} \sqrt{5}} = U\Sigma V^T$.

- 13** The homemade proof depends on this step: If Λ is diagonal and $\Sigma\Lambda = \Lambda\Sigma$ then Σ is also diagonal. That step fails when $\Lambda = I$ because $\Sigma I = I\Sigma$ for all Σ . The step fails unless the numbers $\lambda_1, \dots, \lambda_n$ are all different (which is usually true—but not always, and we want a proof that always works).

Note: If $\lambda_1 \neq \lambda_2$ then comparing the (1, 2) entries of $\Sigma\Lambda$ and $\Lambda\Sigma$ gives $\lambda_2\sigma_{12} = \lambda_1\sigma_{12}$ which forces $\sigma_{12} = 0$. Similarly, all the other off-diagonal σ 's will be zero. Repeated eigenvalues $\lambda_1 = \lambda_2$ or singular values always bring extra steps.

- 14** For a 2 by 3 matrix $A = U\Sigma V^T$, U has 1 parameter (angle θ) and Σ has 2 parameters (σ_1 and σ_2) and V has 3 parameters (3 angles like roll, pitch, and yaw for an aircraft in 3D flight). Total 6 parameters in $U\Sigma V^T$ agrees with 6 in the 2 by 3 matrix A .

- 15** For 3 by 3 matrices, U and Σ and V have 3 parameters each. For 4 by 4, Σ has 4 singular values and U and V involve 6 angles each: $6 + 4 + 6 = 16$ parameters in A . (See also the last Appendix.)
- 16** 4 numbers give a direction in \mathbf{R}^5 . A unit vector orthogonal to that direction has 3 parameters. The remaining columns of Q have 2, 1, 0 parameters (not counting $+/-$ decisions). Total $4 + 3 + 2 + 1 + 0 = 10$ parameters in Q .
- 17** If $A^T A \mathbf{v} = \lambda \mathbf{v}$ with $\lambda \neq 0$, multiply by A : $(AA^T)A \mathbf{v} = \lambda A \mathbf{v}$ with eigenvector $A \mathbf{v}$.
- 18** $A = U \Sigma V^T$ gives $A^{-1} = V \Sigma^{-1} U^T$ when A is invertible. The singular values of $A^T A$ are $\sigma_1^2, \dots, \sigma_r^2$ (squares of singular values of A).
- 19** (Correction to 1st printing: Change S to A : *not symmetric!*) If A has orthogonal columns of lengths 2, 3, 4 then $A^T A = \text{diag}(4, 9, 16)$ and $\Sigma = \text{diag}(2, 3, 4)$. We can choose $V = \text{identity matrix}$ and $U = A \Sigma^{-1}$ has orthogonal *unit vectors*: the original columns divided by 2, 3, 4.
- 21** We know that $AA^T A = (U \Sigma V^T)(V \Sigma^T U^T)(U \Sigma V^T) = U(\Sigma \Sigma^T \Sigma) V^T$. So the singular values from $\Sigma \Sigma^T \Sigma$ are σ_1^3 to σ_r^3 .
- 22** To go from the reduced form $AV_r = U_r \Sigma_r$ to $A = U_r \Sigma_r V_r^T$, we cannot just multiply both sides by V_r^T (Since V_r only has r columns and rank r , possibly a small number, and then $V_r V_r^T$ is not the identity matrix). But the result $A = U_r \Sigma_r V_r^T$ is still correct, since both sides give the zero vector when they multiply the basis vectors $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ in the nullspace of A .
- 23** This problem is solved in the final appendix of the book. Note for $r = 1$ those rank-one matrices have $m + n - 1$ free parameters: vectors \mathbf{u}_1 and \mathbf{v}_1 have $m + n$ parameters but there is freedom to make one of them a unit vector: $A = (\mathbf{u}_1 / \|\mathbf{u}_1\|)(\|\mathbf{u}_1\| \mathbf{v}_1^T)$.

Problem Set I.9, page 80

- 1 The singular values of $A - A_k$ are $\sigma_{k+1} \geq \sigma_{k+2} \geq \dots \geq \sigma_r$ (the smallest $r - k$ singular values of A).

2 The closest rank 1 approximations are $A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ $A = \begin{bmatrix} 0 & 3 \\ 0 & 0 \end{bmatrix}$ $A = \frac{3}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

- 3 Since this A is orthogonal, its singular values are $\sigma_1 = \sigma_2 = 1$. So we cannot reduce its spectral norm σ_{\max} by subtracting a rank-one matrix. On the other hand, we can reduce its Frobenius norm from $\|A\|_F = \sqrt{2}$ to $\|A - \mathbf{u}_1 \sigma_1 \mathbf{v}_1^T\|_F = \sqrt{1}$.

- 4 $A - A_1 = \frac{1}{2} \begin{bmatrix} 3 & -3 \\ -1 & 1 \end{bmatrix}$ has $\|A - A_1\|_{\infty} = \max \text{ row sum} = 3$. But in this “ ∞ norm” (which is not determined by the singular values) we can find a rank-one matrix B that is closer to A than A_1 is.

$$B = \begin{bmatrix} 1 & .75 \\ 4 & 3 \end{bmatrix} \text{ has } A - B = \begin{bmatrix} 2 & -.75 \\ 0 & 2 \end{bmatrix} \text{ and } \|A - B\|_{\infty} = 2.75.$$

- 5 If $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ then $QA = \begin{bmatrix} \cos \theta & 0 \\ \sin \theta & 0 \end{bmatrix}$. Those matrices have $\|A\|_{\infty} = 1$ different from $\|QA\|_{\infty} = |\cos \theta|$.

- 6 $S = Q\Lambda Q^T = \lambda_1 \mathbf{q}_1 \mathbf{q}_1^T + \dots$ is the eigenvalue decomposition and *also the singular value decomposition* of S . So the Eckart-Young theorem applied to $\lambda_1 \mathbf{q}_1 \mathbf{q}_1^T$ is the nearest rank-one matrix.

- 7 Express $E = \|A - CR\|_F^2$ as $E = \sum_{i,j} (A_{ij} - \sum_k C_{ik} R_{kj})^2$. Take the derivative with respect to each particular C_{IK} .

$$\frac{\partial E}{\partial C_{IK}} = 2 \sum_j (A_{Ij} - C_{IK} R_{Kj}) R_{Kj}$$

The (1, 1) entry of $A - CR$ is $a_{11} - c_{11}r_{11} - c_{12}r_{21}$. The (1, 1) entry of $A - (C + \Delta C)R$ is $a_{11} - c_{11}r_{11} - c_{12}r_{21} - \Delta c_{11}r_{11} - \Delta c_{12}r_{21}$. **TO COMPLETE**

Squaring and subtracting, the leading terms (first-order) are $2(a_{11} - c_{11}r_{11} - c_{12}r_{21})(\Delta c_{11}r_{11} + \Delta c_{12}r_{12})$.

- 8** $\|A - A_1\|_2 = \sigma_2(A)$ and $\|A - A_2\|_2 = \sigma_3(A)$. (The 2-norm for a matrix is its largest singular value.) So those norms are equal when $\sigma_2 = \sigma_3$.
- 9** Our matrix has 1's below the parabola $y = 1 - x^2$ and 0's above that parabola. The parabola has slope $dy/dx = -2x = -1$ where $x = \frac{1}{2}$ and $y = \frac{3}{4}$. Remove the rectangle (filled with 1's and therefore rank = 1) below $y = \frac{3}{4}$ and to the left of $x = \frac{1}{2}$. Above that rectangle, between $y = \frac{3}{4}$ and $y = 1$, the rows of A are independent. Beyond that rectangle, between $x = \frac{1}{2}$ and $x = 1$, the columns of A are independent. Since $\frac{1}{4} + \frac{1}{2} = \frac{3}{4}$, the rank of A is approximately $\frac{3}{4}N$.
- 10** A is invertible so $A^{-1} = V\Sigma^{-1}U^T$ has singular values $1/\sigma_1$ and $1/\sigma_2$. Then $\|A^{-1}\|_2 = \max \text{ singular value} = 1/\sigma_2$. And $\|A^{-1}\|_F^2 = (1/\sigma_1)^2 + (1/\sigma_2)^2$.

Problem Set I.10, page 87

$$\mathbf{1} \quad H = M^{-1/2} S M^{-1/2} = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix} \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix} = \begin{bmatrix} 5 & 4/2 \\ 4/2 & 5/4 \end{bmatrix}$$

$$\det(S - \lambda M) = \det \begin{bmatrix} 5 - \lambda & 4 \\ 4 & 5 - 4\lambda \end{bmatrix} = 4\lambda^2 - 25\lambda + 9 = 0$$

$$\det(H - \lambda I) = \det \begin{bmatrix} 5 - \lambda & 2 \\ 2 & \frac{5}{4} - \lambda \end{bmatrix} = \lambda^2 - \frac{25}{4}\lambda + \frac{9}{4} = 0$$

$$\text{By the quadratic formula, } d = \frac{25 \pm \sqrt{25^2 - 144}}{8} = \frac{25 \pm \sqrt{481}}{8}$$

The first equation agrees with the second equation (times 4). The eigenvectors will be too complicated for convenient computation by hand.

Problem Set I.11, page 96

$$\mathbf{1} \quad \|\mathbf{v}\|_2^2 = v_1^2 + \cdots + v_n^2 \leq (\max|v_i|) (|v_1| + \cdots + |v_n|) = \|\mathbf{v}\|_\infty \|\mathbf{v}\|_1$$

$\mathbf{2}$ (Length)² is never negative. We have to simplify that (length)²:

$$\left(\mathbf{v} - \frac{\mathbf{v}^T \mathbf{w}}{\mathbf{w}^T \mathbf{w}} \mathbf{w}\right)^T \left(\mathbf{v} - \frac{\mathbf{v}^T \mathbf{w}}{\mathbf{w}^T \mathbf{w}} \mathbf{w}\right) = \mathbf{v}^T \mathbf{v} - 2 \frac{(\mathbf{v}^T \mathbf{w})^2}{\mathbf{w}^T \mathbf{w}} + \frac{(\mathbf{v}^T \mathbf{w})^2}{\mathbf{w}^T \mathbf{w}} = \mathbf{v}^T \mathbf{v} - \frac{(\mathbf{v}^T \mathbf{w})^2}{\mathbf{w}^T \mathbf{w}} \geq 0.$$

Multiply by $\mathbf{w}^T \mathbf{w}$.

$$\mathbf{3} \quad \|\mathbf{v}\|_2^2 = v_1^2 + \cdots + v_n^2 \leq n \max|v_i|^2 \text{ so } \|\mathbf{v}\|_2 \leq \sqrt{n} \max|v_i|.$$

For the second part, choose $\mathbf{w} = (1, 1, \dots, 1)$ and use Cauchy-Schwarz:

$$\|\mathbf{v}\|_1 = |v_1|w_1 + \cdots + |v_n|w_n \leq \|\mathbf{v}\|_2 \|\mathbf{w}\|_2 = \sqrt{n} \|\mathbf{v}\|_2$$

$\mathbf{4}$ For $p = 1$ and $q = \infty$, Hölder's inequality says that

$$|\mathbf{v}^T \mathbf{w}| \leq \|\mathbf{v}\|_1 \|\mathbf{w}\|_\infty = (|v_1| + \cdots + |v_n|) \max|w_i|$$

Problem Set I.12, page 109

- 1 The v_1 derivative of $(a - v_1 u_1)^2 + (b - v_1 u_2)^2$ is $-2u_1(a - v_1 u_1) - 2u_2(b - v_1 u_2) = 0$.
Dividing by 2 gives $(u_1^2 + u_2^2)v_1 = u_1 a + u_2 b$. In II.2, this will be the normal equation for the best solution v_1 to the 1D least squares problem $\mathbf{u}v_1 = \mathbf{a}_1$.
- 2 Same problem as 1, stated in vector notation.
- 3 This is the same question but now for the second component v_2 . Together with 1, the combined problem is to find the minimizing numbers (v_1, v_2) for $\|\mathbf{a} - \mathbf{v}\mathbf{u}\|^2$ when \mathbf{u} is fixed.
- 4 The combined problem when U is fixed is to choose V to minimize $\|A - UV\|_F^2$. The best V solves $(U^T U)V = U^T A$.
- 5 This alternating algorithm is important! Here the matrices are small and convergence can be tested and seen computationally.
- 6 Rank 1 requires $A = \mathbf{u}\mathbf{u}^T$. All columns of A must be multiples of one nonzero column. (Then all rows will automatically be multiples of one nonzero row.)
- 7 For the fibers of T in each of the three directions, all slices must be in multiples of one nonzero fiber. (**Question**: If this holds in two directions, does it automatically hold in the third direction?)
- 9 (a) The sum of all row sums must equal the sum of all column sums.
(b) In each separate direction, add the totals for all slices in that direction. For each direction, the sum of those totals must be the total sum of all entries in the tensor.

Problem Set II.2, page 135

- 1 The step from $A^T A \mathbf{x} = \mathbf{0}$ to $A \mathbf{x} = \mathbf{0}$ is proved in the problem statement. The opposite statement (if $A \mathbf{x} = \mathbf{0}$ then $A^T A \mathbf{x} = \mathbf{0}$) is the easier direction. So $\mathbf{N}(A^T A) = \mathbf{N}(A)$. Orthogonal to that subspace is the row space of $A^T A = \text{row space of } A$.
- 2 The link from $A = U \Sigma V^T$ to $A^+ = V \Sigma^+ U^T$ shows that A and A^+ have the same number (the rank r) of nonzero singular values. If A is square and $A \mathbf{x} = \lambda \mathbf{x}$ with $\lambda \neq 0$, then $A^+ \mathbf{x} = \frac{1}{\lambda} \mathbf{x}$. Eigenvectors are the same for A and A^+ , eigenvalues are λ and $1/\lambda$ (except that $\lambda = 0$ for A produces $\lambda = 0$ for A^+ !).
- 3 Note that $(\mathbf{v}_i \mathbf{u}_i^T / \sigma_i) (\sigma_j \mathbf{u}_j \mathbf{v}_j^T)$ is zero if $i \neq j$ (because $\mathbf{u}_i^T \mathbf{u}_j = 0$) and it is $\mathbf{v}_i \mathbf{v}_i^T$ if $i = j$. Then the product $(\sum \mathbf{v}_i \mathbf{u}_i^T / \sigma_i) (\sum \sigma_j \mathbf{u}_j \mathbf{v}_j^T)$ just adds up the results $\mathbf{v}_i \mathbf{v}_i^T$ to get the matrix $V V^T = I$.

Problems 12–22 use four data points $\mathbf{b} = (0, 8, 8, 20)$ to bring out the key ideas.




Figure 1: **Problems 12–22:** The closest line $C + Dt$ matches $C \mathbf{a}_1 + D \mathbf{a}_2$ in \mathbf{R}^4 .

- 12 With $\mathbf{b} = 0, 8, 8, 20$ at $t = 0, 1, 3, 4$, set up and solve the normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$. For the best straight line in Figure 1, find its four heights p_i and four errors e_i . What is the minimum value $E = e_1^2 + e_2^2 + e_3^2 + e_4^2$?

- 13** (Line $C + Dt$ does go through p 's) With $b = 0, 8, 8, 20$ at times $t = 0, 1, 3, 4$, write down the four equations $A\mathbf{x} = \mathbf{b}$ (unsolvable). Change the measurements to $p = 1, 5, 13, 17$ and find an exact solution to $A\hat{\mathbf{x}} = \mathbf{p}$.
- 14** Check that $\mathbf{e} = \mathbf{b} - \mathbf{p} = (-1, 3, -5, 3)$ is perpendicular to both columns of the same matrix A . What is the shortest distance $\|\mathbf{e}\|$ from \mathbf{b} to the column space of A ?
- 15** (By calculus) Write down $E = \|A\mathbf{x} - \mathbf{b}\|^2$ as a sum of four squares—the last one is $(C + 4D - 20)^2$. Find the derivative equations $\partial E/\partial C = 0$ and $\partial E/\partial D = 0$. Divide by 2 to obtain the normal equations $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$.
- 16** Find the height C of the best *horizontal line* to fit $\mathbf{b} = (0, 8, 8, 20)$. An exact fit would solve the unsolvable equations $C = 0, C = 8, C = 8, C = 20$. Find the 4 by 1 matrix A in these equations and solve $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$. Draw the horizontal line at height $\hat{x} = C$ and the four errors in \mathbf{e} .
- 17** Project $\mathbf{b} = (0, 8, 8, 20)$ onto the line through $\mathbf{a} = (1, 1, 1, 1)$. Find $\hat{\mathbf{x}} = \mathbf{a}^T \mathbf{b} / \mathbf{a}^T \mathbf{a}$ and the projection $\mathbf{p} = \hat{\mathbf{x}} \mathbf{a}$. Check that $\mathbf{e} = \mathbf{b} - \mathbf{p}$ is perpendicular to \mathbf{a} , and find the shortest distance $\|\mathbf{e}\|$ from \mathbf{b} to the line through \mathbf{a} .
- 18** Find the closest line $b = Dt$, *through the origin*, to the same four points. An exact fit would solve $D \cdot 0 = 0, D \cdot 1 = 8, D \cdot 3 = 8, D \cdot 4 = 20$. Find the 4 by 1 matrix and solve $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$. Redraw Figure 1a showing the best line $b = Dt$ and the \mathbf{e} 's.
- 19** Project $\mathbf{b} = (0, 8, 8, 20)$ onto the line through $\mathbf{a} = (0, 1, 3, 4)$. Find $\hat{\mathbf{x}} = D$ and $\mathbf{p} = \hat{\mathbf{x}} \mathbf{a}$. The best C in Problems 16–17 and the best D in Problems 18–19 do *not* agree with the best (C, D) in Problems 12–15. That is because $(1, 1, 1, 1)$ and $(0, 1, 3, 4)$ are _____ perpendicular.
- 20** For the closest parabola $b = C + Dt + Et^2$ to the same four points, write down the unsolvable equations $A\mathbf{x} = \mathbf{b}$ in three unknowns $\mathbf{x} = (C, D, E)$. Set up the three normal equations $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (solution not required). In Figure 1a you are now fitting a parabola to 4 points—what is happening in Figure 1b?

- 21** For the closest cubic $b = C + Dt + Et^2 + Ft^3$ to the same four points, write down the four equations $A\mathbf{x} = \mathbf{b}$. Solve them by elimination. In Figure 1a this cubic now goes exactly through the points. What are \mathbf{p} and \mathbf{e} ?
- 22** The average of the four times is $\hat{t} = \frac{1}{4}(0 + 1 + 3 + 4) = 2$. The average of the four b 's is $\hat{b} = \frac{1}{4}(0 + 8 + 8 + 20) = 9$.
- (a) Verify that the best line goes through the center point $(\hat{t}, \hat{b}) = (2, 9)$.
- (b) Explain why $C + D\hat{t} = \hat{b}$ comes from the first equation in $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$.

Problem Set IV.1, page 212

- 1 Warning:** References to the complex dot product should be removed. This is just multiplication of the matrices $F\Omega$. Off the diagonal of $F\Omega$ we have $i \neq j$ and the sum S has powers of $w^i \omega^j = w^{i-j}$.

$$S = 1 + w^{i-j} + w^{2(i-j)} + \dots = 0 \text{ in equation (5) when } i \neq j.$$

- 2** If $M = N/2$ then $(w_N)^M = e^{2\pi i M/N} = e^{\pi i} = -1$.

$$\mathbf{3} \quad F_3 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & w & w^2 \\ 1 & w^2 & w^4 \end{bmatrix} \quad \Omega_3 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega^4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & w^2 & w \\ 1 & w^4 & w^2 \end{bmatrix} \quad \begin{array}{l} \text{because } w = \omega^2 \\ \text{when } N = 3 \end{array}$$

The permutation matrix to exchange columns in $\Omega_3 = F_3 P$ is $P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$

Notice that multiplying by P on the right exchanges *columns*.

$$\mathbf{4} \quad C = \frac{1}{N} F^{-1} \mathbf{f} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & \omega & \omega^2 & \omega^3 \\ 1 & \omega^2 & \omega^4 & \omega^6 \\ 1 & \omega^3 & \omega^6 & \omega^9 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 \\ -i \\ (-i)^2 \\ (-i)^3 \end{bmatrix}$$

$$\mathbf{5} \quad F_6 = \begin{bmatrix} I & D \\ I & -D \end{bmatrix} \begin{bmatrix} F_3 & 0 \\ 0 & F_3 \end{bmatrix} \begin{bmatrix} P \end{bmatrix} \text{ with } D = \begin{bmatrix} 1 \\ w \\ w^2 \end{bmatrix} \text{ and } w = e^{2\pi i/6}$$

The even-odd permutation matrix is $P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

6 $N = 6$ means that $w^3 = e^{2\pi i 3/6} = e^{\pi i} = -1$. Then $1 + w^3 = 0$ and $w + w^4 = 0$ and $w^2 + w^5 = 0$.

7 $2\pi a_0 = \int_{-\pi}^{\pi} f(x) dx = \int_{-\pi/2}^{\pi/2} 1 dx = \pi$ so $a_0 =$ **(the average of $f(x)$)**.

$$a_1 \int_{-\pi}^{\pi} \cos^2 x dx = \int_{-\pi}^{\pi} \cos x f(x) dx = \int_{-\pi/2}^{\pi/2} \cos x dx = \sin \frac{\pi}{2} - \sin \left(-\frac{\pi}{2}\right) = 2$$

so $a_1 = 2/(\pi/2) = \frac{\pi}{4}$.

8 If $A = Q$ then the rank-one pieces have $(\mathbf{q}_i \mathbf{q}_i^T)(\mathbf{q}_j \mathbf{q}_j^T) = 0$ since $\mathbf{q}_i^T \mathbf{q}_j = 0$ for $i \neq j$.

9 The vector \mathbf{x} is $\frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$ and $\mathbf{y} = \frac{1}{4} \begin{bmatrix} 1 \\ \omega^3 \\ \omega^6 \\ \omega^9 \end{bmatrix}$ with $\omega = -i$.

Problem Set IV.2, page 220

1 $(2, 1, 3) * (3, 1, 2) = (6, 5, 14, 5, 6)$ Not cyclic

$$(2, 1, 3) \otimes (3, 1, 2) = (6 + 5, 5 + 6, 14) = (11, 11, 14)$$

Check by adding coefficients of c , d , and $c \otimes b$: $(6)(6) = (36)$. See Problem 3.

2 The question asks for a direct proof by comparing all 9 terms on both sides.

$$\text{Left } (c_0d_0 + c_1d_{-1} + c_2d_{-2}) + w^k(c_0d_1 + c_1d_0 + c_2d_{-1}) + w^{2k}(c_0d_2 + c_1d_1 + c_2d_0)$$

$$\text{Right } (c_0 + w^k c_1 + w^{2k} c_2)(d_0 + w^k d_1 + w^{2k} d_2). \text{ USE } w^3 = 1.$$

3 In $c * d$, every number c_i multiplies every number d_j . So when we add up all terms, we get $\sum c_i$ times $\sum d_j$.

4 $Cq_k = \lambda_k(C)$ times q_k and $Dq_k = \lambda_k(D)$ times q_k . Therefore

$$CDq_k = C(\lambda_k(D)q_k) = \lambda_k(C)\lambda_k(D)q_k \text{ and similarly for } DCq_k.$$

5 This 4×4 circulant matrix C is all ones (rank 1). So it has 3 zero eigenvalues and $\lambda_1 = \text{trace of } C = 4$. Those numbers 4, 0, 0, 0 are exactly the components of Fc for $c = (1, 1, 1, 1)$ because the 3 last rows of F_4 add to zero. The sum $1 + z + z^2 + z^3 = 0$ for $z = i, z = i^2$, and $z = i^3$ (i and -1 and $-i$).

6 The “frequency response” uses the angle θ in $C(e^{i\theta}) = \sum c_j e^{ij\theta}$. At the special angles $\theta = 2\pi/N, 4\pi/N, \dots, 2\pi N/N$, those numbers $e^{ij\theta}$ are exactly $w, w^2, \dots, w^N = 1$. Then the sums of $c_j e^{ij\theta}$ are the sums of $c_j w^j$. Those sums are the components of Fc , which are also the eigenvalues of C .

So Problem 6 says: C is invertible when all its eigenvalues are not zero. True!

7 $c \otimes d = e$ means that the cyclic convolution matrices (*circulants*) have $CD = E$. Their eigenvalues have $\lambda_k(C)\lambda_k(D) = \lambda_k(E)$. Suppose we know C and E . Then the eigenvalue $\lambda_k(D)$ is $\lambda_k(E)$ divided by $\lambda_k(C)$. This tells us all the eigenvalues of D , which are the components of Fd . By inverting that DFT matrix we learn the components of d , which tell us the matrix D .

- 8** This problem uses the Schwarz (or Cauchy-Schwarz) inequality $|\mathbf{f}^T \mathbf{g}| \leq \|\mathbf{f}\| \|\mathbf{g}\|$. The vectors \mathbf{f} and \mathbf{g} are \mathbf{c} and $S^n \mathbf{c}$, which have the same norm because S is just a shift. So $\|S^n \mathbf{c}\| = \|\mathbf{c}\|$ (in other words $\|\mathbf{f}\| = \|\mathbf{g}\|$) and the inequality says that $\mathbf{f}^T \mathbf{g} = \mathbf{c}^T S^n \mathbf{c}$ is not greater than $\|\mathbf{c}\| \|S^n \mathbf{c}\| = \|\mathbf{c}\|^2 = \mathbf{c}^T \mathbf{c}$.

This is the zeroth component of the autocorrelation of \mathbf{c} .